

Varieties of representations of groups and varieties of associative algebras

July 21, 2009

E. Aladova^{a,1}, B. Plotkin^b

^a Department of Mathematics, Bar-Ilan University,
52900, Ramat Gan, Israel

E-mail address: aladovael@mail.ru

^b Department of Mathematics, Hebrew University of Jerusalem, 91904, Jerusalem,
Israel

E-mail address: plotkin@macs.biu.ac.il

Abstract

This paper is tightly connected with the book [25]. Since this book had not been translated from Russian into English we give here the brief review of the basic definitions and results from [25]. We prove also new results of the same spirit. They are related to dimension subgroups, varieties of representations of groups and varieties of associative algebras. The main emphasis is put on the varieties of representations of groups induced by the varieties of associative algebras. We provide the reader with the list of some open problems.

Contents

1	Introduction	2
2	Varieties of representations of groups. General view	2
2.1	Category and variety Rep-K	2
2.2	Varieties of representations as subvarieties of the Rep-K	6
2.3	Algebra of varieties	6
3	Varieties of representations of associative algebras	9
4	Varieties of representation of groups associated with varieties of associative algebras	12
5	General view on dimension subgroups	18
5.1	Dimension subgroup for a variety of representations of groups . .	18
5.2	Linearization of groups and <i>PI</i> -groups	20
5.3	Dimension subgroup for varieties of representations of groups and for varieties of associative algebras	22

¹ The author is supported by the Israel Science Foundation grant number 1178/06.

1 Introduction

It is well known that there are close connections between varieties of groups and varieties of representations of groups, between varieties of algebras and varieties of representations of algebras. The main goal of this paper is to consider some relations between varieties of representations of groups and varieties of associative algebras. In particular, we consider polynomial varieties of representations of groups which are related to varieties of associative algebras.

The set of all varieties of representations of groups is a free semigroup under multiplication of varieties [24]. All polynomial varieties of representations of groups form a subsemigroup in this semigroup. So we come up with the problem: is the semigroup of all polynomial varieties of representations of groups free? To solve this problem we are looking for some examples of polynomial and non-polynomial varieties. For instance, the variety generated by the representation $(K^n, T_n(K))$ is a polynomial variety, while the variety generated by the representation $(K^n, UT_n(K))$, is a non-polynomial variety. Here K is a field, $T_n(K)$ is the group of triangular matrices and $UT_n(K)$ is the group of unitriangular matrices over K which act naturally on the space K^n .

The other our interest is concerned with dimension subgroup. There exist a lot of problems around classical dimension subgroups (see [7], [15], [16], [19], [26] and references therein). We define a concept of the dimension subgroup for varieties of representations of groups. Here we consider the dimension subgroup of a group G as the intersection of kernels of all representations of the group G in the given variety of representations of groups. In particular, we especially distinguish the case of polynomial varieties of representations of groups. We also give a concept of the dimension subgroup for varieties of associative algebras, and consider some problems concerning these subgroups.

A few words about the structure of the paper. In Section 2 we give basic definitions: we define the category **Rep- K** which is the category of all representations of groups over a fixed ring K ; the variety **Rep- K** , and consider some operations on the varieties, in particular, triangular product of varieties of representations of groups. In Section 3 we recall known results about varieties of representations of algebras and varieties of algebras from the book [25]. Some of these results which are very important for us we give with the proofs. In Section 4 we study relations between varieties of representations of groups and varieties of associative algebras. Here we deal with the following problem: to find connection between interesting varieties of representations of groups and some well-known varieties of associative algebras. In the last section we give concepts of the dimension subgroup for variety of representations of groups and variety of associative algebras, and compare two approaches to the notion of the dimension subgroup.

2 Varieties of representations of groups. General view

2.1 Category and variety Rep- K

1.1.1 Category Rep- K . Let K be a commutative and associative ring with unit. Consider pairs (V, G) where V is a K -module and G is a group acting on

the module V . In other words, for each (V, G) there is a function from $V \times G$ to V defined by $(v, g) \rightarrow v \circ g$ satisfying the following conditions:

1. $v_1 \circ (g_1 g_2) = (v_1 \circ g_1) \circ g_2$;
2. $v_1 \circ 1 = v_1$, where 1 is the unit of the group G ;
3. $(v_1 + v_2) \circ g_1 = v_1 \circ g_1 + v_2 \circ g_1$;
4. $(\lambda v_1) \circ g_1 = \lambda(v_1 \circ g_1)$,

where $v_1, v_2 \in V$, $g_1, g_2 \in G$, $\lambda \in K$. The pair (V, G) can be viewed as two-sorted algebra-representation over the ring K (for terminology see books [20], [25]).

On the other hand, if we have a group G acting on a K -module V then there is a representation $\rho : G \rightarrow \text{Aut } V$ of the group G in the K -module V defined as follows $g^\rho(v) = v \circ g$, where $g \in G$, $v \in V$, g^ρ is the image of g under the homomorphism ρ .

Let (V, G) be a representation. Then V can be viewed as a G -module. Moreover, since the group G acts on the K -module V then the group algebra KG also acts on V and, consequently, V is a KG -module.

The class of all pairs (V, G) over a fixed ring K forms the category **Rep- K** . Objects of this category are representations (V, G) , morphisms are homomorphisms $\mu = (\alpha, \beta)$ of representations defined as follows:

Definition 2.1 *A homomorphism of a representation (V_1, G_1) to a representation (V_2, G_2) is a pair of homomorphisms*

$$\mu = (\alpha, \beta) : (V_1, G_1) \rightarrow (V_2, G_2),$$

where $\alpha : V_1 \rightarrow V_2$ is a homomorphism of K -modules, $\beta : G_1 \rightarrow G_2$ is a homomorphism of groups and the following condition holds

$$(v \circ g)^\alpha = v^\alpha \circ g^\beta.$$

Definition 2.2 *A representation (U, H) is a subrepresentation of a representation (V, G) if H is a subgroup of the group G , U is a K -submodule of V and the restriction of the action G on V coincides with the action H on U .*

Along with the subrepresentations we consider a concept of overrepresentation. The defined below induced representations is a particular case of this concept.

Let H be a subgroup of a group G , and let (U, H) be a representation of the group H . Consider the following category: objects of this category are H -homomorphisms $\varphi : U \rightarrow V$ of the H -module U to G -modules V , morphisms of an object $\varphi_1 : U \rightarrow V_1$ to an object $\varphi_2 : U \rightarrow V_2$ are G -homomorphisms $\psi : V_1 \rightarrow V_2$ subject to the following commutative diagram

$$\begin{array}{ccc} & \varphi_1 \nearrow & V_1 \\ U & & \downarrow \psi \\ & \varphi_2 \searrow & V_2 \end{array}$$

Let $\tau : U \rightarrow V$ be an initial object of this category. A representation (V, G) is called induced by the representation (U, H) with the given embedding of the

group H to the group G . Note that U is a right KH -module and KG is a left KG -module. Then we can consider the module $V = U \otimes_{KH} KG$ and the representation (V, G) is the representation induced by the representation (U, H) .

There are products and coproducts of objects in the category **Rep- K** [11].

Definition 2.3 *A representation (V, G) is called the Cartesian product of representations (V_i, G_i) if V is a Cartesian product of the K -modules V_i , G is a Cartesian product of the groups G_i and for all $v \in V$ and all $g \in G$ we have $(v \circ g)(i) = v(i) \circ g(i)$.*

The Cartesian product (V, G) is a product of the objects (V_i, G_i) in the category **Rep- K** .

Definition 2.4 *A representation (V, G) is called the free product of the representations (V_i, G_i) if G is the free product of the groups G_i , V is the direct sum of G -submodules U_i such that representations (U_i, G) are induced by the representations (V_i, G_i) with the natural embedding $G_i \rightarrow G$.*

The free product (V, G) is a coproduct of the objects (V_i, G_i) in the category **Rep- K** .

Now we describe a free objects in the category **Rep- K** . Let (V, G) be an arbitrary representation. Let Y be a subset of G and let X be a subset of V . We say that the pair of the sets $\{X, Y\}$ generates the representation (V, G) if the group G is generated by the set Y and the module V is generated by the set X as KG -module.

Definition 2.5 *A representation (W, F) is called a free representation over a pair of the sets $\{X, Y\}$ if for every representation (V, G) any pair of the functions $\mu_1 : Y \rightarrow G$ and $\mu_2 : X \rightarrow V$ can be extended uniquely up to the homomorphism of the representations $(W, F) \rightarrow (V, G)$.*

1.1.2 Variety Rep- K . Along with the category **Rep- K** we consider variety of representations of groups **Rep- K** . This variety consists of all two-sorted algebras-representations. Identities of the variety **Rep- K** are identities of groups, identities of K -modules and identities of action written earlier.

Describe the free representations in the variety **Rep- K** . Let $\{X, Y\}$ be a pair of sets. Let $F = F(Y)$ be the free group over the set Y , $W = W(X, Y)$ be the free KF -module over the set X . Moreover, we have

$$W(X, Y) = X \cdot KF = \bigoplus_{x \in X} x \cdot KF,$$

and each element $w \in W$ have the form:

$$w = x_1 u_1 + \dots + x_n u_n,$$

where $u_i = u_i(y_1, \dots, y_m) \in KF$, $x_i \in X$. Define an action of the group F on the module W as follows:

$$w \circ f = (x_1 u_1 + \dots + x_n u_n) \circ f = x_1 (u_1 f) + \dots + x_n (u_n f),$$

where $w \in W$, $f \in F$, $u_i f \in KF$.

Definition 2.6 A representation (W, F) with the action $w \circ f$ above is the free representation over $\{X, Y\}$.

So, we can speak about two types of identities of representations:

$$w \equiv 0 \text{ is an identity of action, } w \in W(X, Y),$$

and

$$f \equiv 1 \text{ is a group identity, } f \in F.$$

In fact we will consider only identities of the action. Moreover, it is sufficient to consider identities of the form

$$x \circ u \equiv 0, \quad u \in KF, \quad x \in X.$$

Let F be a free group of countable rank with free generators $y_1, y_2, \dots$, and let KF be a group algebra of F . Let $u(y_1, \dots, y_n)$ be an element of KF .

Definition 2.7 We say that a representation (V, G) satisfies an identity $x \circ u(y_1, \dots, y_n) \equiv 0$ if for all $v \in V$ and all $g_i \in G$ we have $v \circ u(g_1, \dots, g_n) = 0$.

Consider some examples of identities of representations of groups.

Examples.

1. A representation (V, G) satisfies the identity

$$x \circ (y - 1) \equiv 0$$

if the group G acts trivially on the module V .

2. A representation (V, G) satisfies the identity

$$x \circ (y_1 - 1)(y_2 - 1) \dots (y_n - 1) \equiv 0$$

if the module V has a G -invariant series of submodules of the length n :

$$0 = V_0 \subset V_1 \subset \dots \subset V_n = V,$$

such that the group G acts trivially on the factors. The representation $(K^n, UT_n(K))$ satisfies this identity. Here $UT_n(K)$ is the group of unitriangular matrices over a field K , and this group acts in a natural way on the module K^n .

3. Let (V, G) be a finite dimension representation over a field K and let $n = \dim V$. It is known that the algebra $\text{End} V$ satisfies the standard identity of Amitsur-Levitzky of the degree $2n$, that is

$$s_{2n} = \sum_{\sigma \in S_{2n}} (-1)^{\text{sgn} \sigma} x_{\sigma(1)} \dots x_{\sigma(2n)} \equiv 0,$$

where S_{2n} is the symmetric group. So all finite dimension representations possess non-trivial identities.

2.2 Varieties of representations as subvarieties of the $\mathbf{Rep}\text{-}K$

In this part of the paper we will speak about varieties of representations which are subvarieties of $\mathbf{Rep}\text{-}K$. This means that varieties in question satisfy a system of identities of the form $x \circ u \equiv 0$. We suppose that the free group $F = F(Y)$ is a group of countable rank.

Let $\mathfrak{X}$ be a class of representations from the variety $\mathbf{Rep}\text{-}K$. Let $I_{\mathfrak{X}}(KF)$ be a set of all $u \in KF$ such that the identity $x \circ u \equiv 0$ holds in every representation from $\mathfrak{X}$. It is easy to see that the set $I_{\mathfrak{X}}(KF)$ is a two-sided ideal of the group algebra KF which is invariant under all endomorphisms of the group F , i.e., $I_{\mathfrak{X}}(KF)$ is a fully invariant ideal of KF . Furthermore there is a one-to-one correspondence between the varieties of representations of groups over K and the fully invariant ideals of KF [25].

Let $\mathfrak{X}$ be a class of representations of groups over K . Denote by $\text{Var}\mathfrak{X}$ the variety of representations generated by the class $\mathfrak{X}$. For varieties of representations of groups we have the following invariant description [25]:

Theorem 2.8 *A class of representations of groups forms a variety of representations of groups if and only if it is closed under subrepresentations, homomorphic images, Cartesian products and saturations.*

Remind that a class of representations of groups $\mathfrak{X}$ is closed under the saturation if the following condition is true: if a representation (V, H) lies in the class $\mathfrak{X}$ then all representations (V, G) such that (V, H) is a right epimorphic image of the representation (V, G) also belong to $\mathfrak{X}$.

2.3 Algebra of varieties

1.3.1 Operations on varieties. Now we consider the algebra of subvarieties of $\mathbf{Rep}\text{-}K$.

Let $\mathfrak{X}_1$ and $\mathfrak{X}_2$ be two subvarieties from $\mathbf{Rep}\text{-}K$.

Definition 2.9 ([25]) *A product of varieties $\mathfrak{X}_1$ and $\mathfrak{X}_2$ is a variety $\mathfrak{X}_1\mathfrak{X}_2$ defined as follows: a representation (V, G) belongs to $\mathfrak{X}_1\mathfrak{X}_2$ if and only if the module V has a G -submodule U such that $(U, G) \in \mathfrak{X}_1$ and $(V/U, G) \in \mathfrak{X}_2$.*

The operation of the multiplication of varieties is associative. Thus we have a semigroup $\mathfrak{M} = \mathfrak{M}(K)$ of varieties of representations of groups.

For example, let $\mathfrak{S}$ be a variety satisfying the identity $x \circ (y - 1) \equiv 0$ and $\mathfrak{S}^n$ be a variety determined by the identity $x \circ (y_1 - 1) \dots (y_n - 1) \equiv 0$. Then the variety $\mathfrak{S}^n$ is n -th degree of the variety $\mathfrak{S}$.

There is the following theorem [24] (see also [25]):

Theorem 2.10 *If K is a field then the semigroup $\mathfrak{M} = \mathfrak{M}(K)$ is a free semigroup.* $\square$

In the proof of Theorem 2.10 the construction of triangular product of representations is used. Define this structure (see [25], [29]). Let (V_1, G_1) and (V_2, G_2) be representations. Define a representation (V, G) as follows

$$(V, G) = (V_1, G_1) \nabla (V_2, G_2),$$

where $V = V_1 \oplus V_2$ is the direct sum of the modules and elements of the acting group G are matrices of the form:

$$g = \begin{pmatrix} g_1 & \varphi \\ 0 & g_2 \end{pmatrix},$$

where $g_1 \in G_1$, $g_2 \in G_2$, $\varphi \in \text{Hom}(V_1, V_2)$.

Let

$$g = \begin{pmatrix} g_1 & \varphi \\ 0 & g_2 \end{pmatrix}, \quad g' = \begin{pmatrix} g'_1 & \varphi' \\ 0 & g'_2 \end{pmatrix}$$

be elements of the group G . The product of these elements is defined as follows:

$$gg' = \begin{pmatrix} g_1 g'_2 & \overline{g_1} \varphi' + \varphi \overline{g'_2} \\ 0 & g_2 g'_2 \end{pmatrix},$$

where $g_i, g'_i \in G_i$, $\varphi, \varphi' \in \text{Hom}(V_1, V_2)$, $\overline{g_1} \in \overline{G_1}$, $\overline{g'_2} \in \overline{G_2}$ and $(V_i, \overline{G_i})$ are faithful representations corresponding to the representations (V_i, G_i) , $i = 1, 2$.

The group G acts on the module V by the rule:

$$(v_1 + v_2) \circ \begin{pmatrix} g_1 & \varphi \\ 0 & g_2 \end{pmatrix} = v_1 \circ g_1 + \varphi(v_1) \circ g_2 + v_2 \circ g_2,$$

where $v_1 \in V_1$, $v_2 \in V_2$.

Note that the submodule V_1 of the module V is invariant under the action of the group G . Moreover the representation (V_1, G) has the same identities as the representation (V_1, G_1) and the representation $(V/V_1, G)$ has the same identities as the representation (V_2, G_2) .

Henceforward K is a field. The basic role in the proof of Theorem 2.10 plays the following proposition.

Proposition 2.11 ([25]) *Over a field we have the following property:*

$$\text{Var}((V_1, G_1) \nabla (V_2, G_2)) = \text{Var}(V_1, G_1) \cdot \text{Var}(V_2, G_2). \quad \square$$

Let Θ_1 and Θ_2 be varieties of groups.

Definition 2.12 *A product of varieties of groups Θ_1 and Θ_2 is a class $\Theta_1 \Theta_2$ defined as follows: a group G belongs to the variety $\Theta_1 \Theta_2$ if it has an invariant subgroup H such that $H \in \Theta_1$ and $G/H \in \Theta_2$.*

Note that the class $\Theta_1 \Theta_2$ is a variety of groups. Let $\mathfrak{M}^g$ be the semigroup of varieties of groups. It is well-known that $\mathfrak{M}^g$ is a free semigroup [18], [28].

Now we assign an action of the semigroup $\mathfrak{M}^g$ on the semigroup $\mathfrak{M}$ of varieties of representations of groups.

Let $\mathfrak{X}$ be a variety of representations of groups and Θ be a variety of groups. Define a class of representations of groups $\mathfrak{X} \times \Theta$ as follows:

Definition 2.13 ([25]) *A representation (V, G) belongs to $\mathfrak{X} \times \Theta$ if the group G has a normal subgroup H such that $(V, H) \in \mathfrak{X}$ and $G/H \in \Theta$.*

Note that the class $\mathfrak{X} \times \Theta$ forms a variety of representations of groups and there is the following

Proposition 2.14 ([25]) *Let $\mathfrak{X}_1$ and $\mathfrak{X}_2$ be varieties of representations of groups and let Θ_1 and Θ_2 be varieties of groups. Then*

1. $\mathfrak{X}_1 \times \Theta_1 \Theta_2 = (\mathfrak{X}_1 \times \Theta_1) \times \Theta_2$,
2. $\mathfrak{X}_1 \mathfrak{X}_2 \times \Theta_1 = (\mathfrak{X}_1 \times \Theta_1)(\mathfrak{X}_2 \times \Theta_1)$. $\square$

The following theorem is a main theorem of this theory.

Theorem 2.15 ([22]) *If K is a field then the semigroup $\mathfrak{M}^g$ acts freely on the semigroup $\mathfrak{M} = \mathfrak{M}(K)$. $\square$*

In particular, this theorem yields the following property: for every variety of representations of groups $\mathfrak{X}$ there is unique decomposition

$$\mathfrak{X} = (\mathfrak{X}_1 \times \Theta_1) \dots (\mathfrak{X}_n \times \Theta_n),$$

where all $\mathfrak{X}_i$ are indecomposable as $(\mathfrak{X}_{i_1} \times \Theta_{i_1})$.

Example. Let $\mathfrak{S}_n$ be the variety determined by the identity $x \circ (y - 1)^n \equiv 0$. The variety $\mathfrak{S}_n$ is indecomposable.

In the proof of Theorem 2.15 the construction of the wreath product of a representation of a group and a group is used.

Let (V, H) be a representation and G be a group. Define

$$(V, H) \text{ wr } G = (V^G, H \text{ wr } G) = (V^G, H^G \rtimes G).$$

Only in special cases we have the following property:

$$\text{Var}((V, G) \text{ wr } G) = \text{Var}(V, G) \times \text{Var } G.$$

1.3.2 Varieties of groups and varieties of representations of groups.

Now we give some remarks about connections between varieties of groups and varieties of representations of groups.

Let Θ be a variety of groups. Define a class of representations of groups $\omega\Theta$ as follows

Definition 2.16 *Let (V, G) be a representation and $(V, \overline{G})$ the corresponding faithful representation. A representation (V, G) lies in $\omega\Theta$ if the group $\overline{G}$ belongs to Θ .*

Note that class $\omega\Theta$ is a variety of representations of groups. It is obvious that $\omega\Theta = \mathfrak{S} \times \Theta$, where $\mathfrak{S}$ is the variety of representations satisfying the identity $x \circ (y - 1) \equiv 0$.

Let $\mathfrak{A}$ be a variety of abelian group, $\mathfrak{A}_m$ be a variety of abelian groups of exponent m . Let K be an integral domain and let K^* be a multiplicative group of K . Then we have two propositions (see [25]):

Proposition 2.17 *If K^* is infinite then the variety of representations of groups $\omega\mathfrak{A}$ is generated by the representation (K, K^*) , where K^* acts on the cyclic module K by multiplication. That is*

$$\omega\mathfrak{A} = \text{Var}(K, K^*).$$

Proposition 2.18 *If the order of K^* is equal m then the variety of representations of groups $\omega\mathfrak{A}_m$ is generated by the representation (K, K^*) .*

Using the proposition 2.14 we have the following fact:

$$(\omega\mathfrak{A})^n = (\mathfrak{S} \times \mathfrak{A})^n = \mathfrak{S}^n \times \mathfrak{A}.$$

The variety $\mathfrak{S}^n$ is generated by the representation $(K^n, UT_n(K))$ (see [5]), and the variety $\mathfrak{S}^n \times \mathfrak{A}$ is generated by the representation $(K^n, T_n(K))$ ([10], see also [25]).

Now define the operator ω' from varieties of groups Θ to varieties of representations of groups $\omega'\Theta$. Let (V, G) be a representation of a group. Consider V as an abelian group and take the semidirect product of groups $V \rtimes G$. We say that the representation (V, G) belongs to the class $\omega'\Theta$ if $V \rtimes G \in \Theta$. The class $\omega'\Theta$ is a variety of representations of groups. Here the symbol $'$ is interrelated with the following fact: the identities of the variety $\omega'\Theta$ are given by the special Fox derivatives used in the theory of formal power series [4]. We have the following conditions:

1. $\omega'(\Theta_1\Theta_2) = ((\omega'\Theta_1) \times \Theta_2) \cdot \omega'\Theta_2$,
2. $\omega'[\Theta_1, \Theta_2] = (\omega\Theta_1 \cdot \omega'\Theta_2) \cap (\omega\Theta_2 \cdot \omega'\Theta_1)$,

where $[\Theta_1, \Theta_2]$ is the commutator of varieties. Second formula confirms the "differential character" of the operator ω' .

In the sequel we will give some other remarks and propositions concerning this theory.

3 Varieties of representations of associative algebras

In this section we consider some definitions and facts concerning varieties of representations of associative algebras. Here we rely on §14 of the book [25] since there is an essential connections of these facts with new results.

Let K be an associative and commutative ring with unit, let A be an associative algebra over K and let V be a K - and A -module. Let (V, A) be a representation of the algebra A . We also can consider the category of representations of associative algebras over a fixed ring K and various algebras A . A variety Σ in this category can be defined as a class of representations of algebras closed under the subrepresentations, homomorphic images, Cartesian products and saturations. These operators are defined in the same way as for representations of groups (see section 2.1, 2.2). Similarly we can define the product of two representations of algebras over a fixed ring K (see section 2.3) and so we have the semigroup of representations of algebras $\mathfrak{M}^a = \mathfrak{M}^a(K)$.

Let Σ be a variety of associative K -algebras. Define a class of representations of algebras $\omega\Sigma$ as follows:

Definition 3.1 *A representation (V, A) belongs to $\omega\Sigma$ if the algebra $\overline{A}$, where $(V, \overline{A})$ is faithful belongs to Σ .*

The class of representations $\omega\Sigma$ is a variety of representations. A variety of representations of algebras of the form $(0, A)$, where 0 is the zero algebra is called a singular variety. Let $\mathfrak{V}$ be a non-singular variety of representations of algebras.

Definition 3.2 *Denote by $\overrightarrow{\mathfrak{V}}$ the class of all algebras admitting a faithful representation in the variety $\mathfrak{V}$.*

We will prove that the class $\overrightarrow{\mathfrak{V}}$ is a variety of algebras. Note that this fact makes difference between the cases of algebras and groups. It is known [25] that a class $\overrightarrow{\mathfrak{X}}$ of all groups admitting a faithful representation in the variety of representations of groups $\mathfrak{X}$ constitutes a quasivariety of groups.

Let A be a K -algebra with unit and J be an ideal of K annihilating the K -module A . We can consider the algebra A as a K/J -algebra.

Let $\mathfrak{V}$ be a variety of representations of algebras over a ring K and let $\mathbb{F}$ be a free K -algebra with unit. Let $I_{\mathfrak{V}}(\mathbb{F}) \triangleleft \mathbb{F}$ be the ideal of identities of the variety $\mathfrak{V}$. So the representation $(\mathbb{F}/I_{\mathfrak{V}}(\mathbb{F}), \mathbb{F})$ is the free representation of the variety $\mathfrak{V}$. Let J be an ideal of K of all $\alpha \in K$ such that α annihilates K -module $\mathbb{F}/I_{\mathfrak{V}}(\mathbb{F})$, i.e., $\alpha \in J$ if $\alpha \cdot 1 \in I_{\mathfrak{V}}(\mathbb{F})$. It is clear that, if (V, A) is a representation from the variety $\mathfrak{V}$ then the ideal J annihilates the module V . Moreover, if A is an algebra from $\overrightarrow{\mathfrak{V}}$ then the ideal J annihilates A . The ideal J is called a K -annihilator of the variety $\mathfrak{V}$.

We have the following

Lemma 3.3 ([25]) *Let $\mathfrak{V}$ be a variety of representations of algebras. Then for every algebra A from the variety $\overrightarrow{\mathfrak{V}}$ the regular representation (A, A) belongs to the variety $\mathfrak{V}$.*

Proof. Since $A \in \overrightarrow{\mathfrak{V}}$ then there exists a faithful representation (V, A) from $\mathfrak{V}$. Let $U = K/J$, where J is the K -annihilator of $\mathfrak{V}$. The representation $(U, 0)$ also belongs to $\mathfrak{V}$ as a subrepresentation of the free representation $(\mathbb{F}/I_{\mathfrak{V}}(\mathbb{F}), \mathbb{F})$. Using the homomorphism $A \rightarrow 0$ we define a zero representation (U, A) and $(U, A) \in \mathfrak{V}$.

Let us consider the A -module $V \oplus U$. The representation $(V \oplus U, A)$ is a faithful representation and it belongs to $\mathfrak{V}$.

Now we consider the representation (A, A) . Let w be an element of $V \oplus U$. Define the map $\mu_w : A \rightarrow V \oplus U$ by the rule $\mu_w(a) = w \circ a$, where $a \in A$. The map μ_w is a homomorphism of K - and A -modules. Let $U_w = \text{Ker } \mu_w$. Then $A/U_w \leq V \oplus U$ and the representation $(A/U_w, A)$ belongs to the variety $\mathfrak{V}$ as a subrepresentation of the representation $(V \oplus U, A) \in \mathfrak{V}$. Since $\mathfrak{V}$ is a variety then the representation $(\prod_w (A/U_w), A)$ belongs to the variety $\mathfrak{V}$. The algebra A acts faithfully on $V \oplus U$ and $\bigcap U_w = \{0\}$. Using the Remak's theorem we have $A \leq \prod_w (A/U_w)$. Thus the representation (A, A) belongs to $\mathfrak{V}$ as a subrepresentation of $(\prod_w (A/U_w), A)$. $\square$

We have the following

Proposition 3.4 ([25]) *Let $\mathfrak{V}$ be a variety of representations of associative K -algebras. The class $\overrightarrow{\mathfrak{V}}$ is a variety of associative algebras.*

Proof. It is necessary to prove that the class $\mathfrak{V}$ closed under subalgebras, Cartesian products and homomorphic images.

It is obvious that first two conditions are hold. We show that the class $\overrightarrow{\mathfrak{V}}$ closed under homomorphic images. Let A be an algebra from the class $\overrightarrow{\mathfrak{V}}$ and let $\mu : A \rightarrow A_1$ be an epimorphism of algebras. The epimorphism μ induces an epimorphism of the representations $\overrightarrow{\mu} : (A, A) \rightarrow (A_1, A_1)$. Since the regular representation (A, A) belongs to $\mathfrak{V}$ then the regular representation (A_1, A_1) also belongs to $\mathfrak{V}$. This representation is faithful, so the algebra A_1 belongs to $\overrightarrow{\mathfrak{V}}$. Thus the class $\overrightarrow{\mathfrak{V}}$ forms a variety of algebras. $\square$

Thus the map $\rightarrow$ is a map from the set of all varieties of representations of algebras over K to the set of all varieties of algebras. This fact is not true for varieties of representations of groups and for varieties of representations of Lie algebras (see [25]). But we can note that always there is the following property $\overrightarrow{\omega\Sigma} = \Sigma$ (where Σ is a variety of groups, algebras or Lie algebras).

Let Σ be a variety of associative algebras over a ring K , let J be an ideal of K annihilating all algebras from the variety Σ . Denote by $\omega_J\Sigma$ an intersection of the variety of representations $\omega\Sigma$ and the variety of all representations (V, A) such that the ideal J annihilates the module V . Then the following proposition holds:

Proposition 3.5 ([25]) *Let $\mathfrak{V}$ be a variety of representation of algebras over a ring K , let J be the K -annihilator of the variety $\mathfrak{V}$. Then*

$$\mathfrak{V} = \omega_J\overrightarrow{\mathfrak{V}}.$$

Proof. It is not difficult to see that $\mathfrak{V} \subseteq \omega_J\overrightarrow{\mathfrak{V}}$. We prove that $\omega_J\overrightarrow{\mathfrak{V}} \subseteq \mathfrak{V}$. Let (V, A) be a representation from $\omega_J\overrightarrow{\mathfrak{V}}$. We consider two cases: V is a cyclic A -module and general case.

Let V be a cyclic A -module and let $(V, \overline{A})$ be the faithful representation corresponding the representation (V, A) . Then $\overline{A} \in \overrightarrow{\mathfrak{V}}$. According to the lemma 3.3 the regular representation $(\overline{A}, \overline{A})$ belongs to $\mathfrak{V}$. The representation $(V, \overline{A})$ is a homomorphic image of the representation $(\overline{A}, \overline{A})$, and then the representation $(V, \overline{A})$ belongs to the variety $\mathfrak{V}$. Since $(V, \overline{A})$ is a right epimorphic image of the representation (V, A) and $(V, \overline{A}) \in \mathfrak{V}$ then (V, A) belongs to $\mathfrak{V}$.

Now let (V, A) be an arbitrary representation from $\omega_J\overrightarrow{\mathfrak{V}}$. Since the A -module V is generated by its cyclic submodules V_i and the representations (V_i, A) belongs to $\mathfrak{V}$ then the representation (V, A) belongs to $\mathfrak{V}$. $\square$

We have the following

Theorem 3.6 ([25]) *Let K be a field. Then there is a one-to-one correspondence between varieties of K -algebras and varieties of representations of algebras over K .*

Proof. From the propositions 3.4 and 3.5 follows that if K is a field then the maps ω and $\rightarrow$ are mutually inverse. So the statement of the theorem is true. $\square$

Now we give some remarks about identities of varieties of representations of algebras.

Let (V, A) be a faithful representation over a ring K from a variety $\mathfrak{V}$, and let $\mathbb{F}$ be the free K -algebra. An element $u \in \mathbb{F}$ is an identity of the faithful representation (V, A) if and only if u is the identity of the algebra A .

Thus, if $\mathfrak{V}$ is a variety of representations of algebras over a ring K and $\mathfrak{V} = \omega_J \Sigma$, where $\Sigma = \overline{\mathfrak{V}}$, then the ideal of identities of the variety $\mathfrak{V}$ is generated by the ideal of identities of the variety Σ and the elements $\lambda \cdot 1$, where $\lambda \in J$.

Note that, if K is a field then the ideal of identities of a variety of representations of algebras $\mathfrak{V}$ coincides with the ideal of identities of the variety of algebras $\Sigma = \overline{\mathfrak{V}}$.

Now let K be a field, let $\mathbb{F}$ be a free associative K -algebra of countable rank and let Σ be a class of associative K -algebras. Take $u_i \in \mathbb{F}$ such that $u_i \equiv 0$ are identities in all algebras from the class Σ . Then the elements u_i form an ideal of the algebra $\mathbb{F}$ closed under all endomorphisms of $\mathbb{F}$. This ideal is called a fully invariant ideal (or T -ideal) and moreover there is one-to-one correspondence between fully invariant ideals of the algebra $\mathbb{F}$ and varieties of associative algebras over a field K .

Let Σ_1 and Σ_2 be varieties of algebras over a field K , let I_1 and I_2 be fully invariant ideals (T -ideals) of the varieties Σ_1 and Σ_2 accordingly. Let consider varieties of representations of algebras $\omega \Sigma_1$ and $\omega \Sigma_2$ and their product $\omega \Sigma_1 \cdot \omega \Sigma_2$. Let $\Sigma = \omega^{-1}(\omega \Sigma_1 \cdot \omega \Sigma_2)$ and let I be an ideal of identities of the variety Σ . It is not difficult to check that $I = I_2 \cdot I_1$.

So the semigroup $\mathfrak{M}^a(K)$ of all varieties of representations of algebras over a given field K is anti-isomorphic to the semigroup of all T -ideals of the free K -algebra $\mathbb{F}$. The well-known theorem of Bergman and Lewin [2] states that the semigroup of all T -ideals of $\mathbb{F}$ is a free semigroup. Note, that this theorem can be proved with a help of the construction of the triangular product (see section 2.3).

4 Varieties of representation of groups associated with varieties of associative algebras

Let K be a field and let Σ be a variety of associative K -algebras with unit. Define a class $\eta \Sigma$ of representations of groups as follows:

Definition 4.1 *A representation (V, G) belongs to $\eta \Sigma$ if the representation (V, KG) lies in the variety $\omega \Sigma$.*

In other words, if $(V, \overline{G})$ is the faithful representation of group $\overline{G}$ corresponding to the representation (V, G) then the linear span of $\overline{G}$ in the algebra $\text{End} V$ is an algebra from the variety Σ .

We have the following

Proposition 4.2 ([25]) *The class $\eta \Sigma$ is a variety of representations of groups.*

Proof. Let (V, G) be a representation from the variety $\eta \Sigma$ and let (U, H) be a subrepresentation of (V, G) . Then (U, KH) is a subrepresentation of (V, KG) . The representation (V, KG) belongs to variety $\omega \Sigma$, so (U, KH) also belongs to $\omega \Sigma$. Hence (U, H) is a representation from the variety $\eta \Sigma$. Thus the class $\eta \Sigma$ is closed under subrepresentations. In a similar way it can be proved that the class $\eta \Sigma$ is closed under homomorphic images.

Now we prove that the class $\eta\Sigma$ is closed under saturations. Let (V, G) be a representation such that the faithful representation $(V, \overline{G})$ belongs to $\eta\Sigma$. Then $(V, K\overline{G})$ lies in $\omega\Sigma$ and the faithful representations for (V, KG) and $(V, K\overline{G})$ are isomorphic. So (V, KG) belongs to $\omega\Sigma$ and the representation (V, G) belongs to $\eta\Sigma$.

It remains to prove that the class $\eta\Sigma$ is closed under Cartesian products. Let (V, G) be a Cartesian product of the representations (V_i, G_i) , $i \in I$, and $(V_i, G_i) \in \eta\Sigma$. Let (V, A) be a Cartesian product of the representations (V_i, KG_i) . For every projection $\pi_i : (V, G) \rightarrow (V_i, G_i)$ we have epimorphism $\nu_i : (V, KG) \rightarrow (V_i, KG_i)$. All epimorphisms ν_i can be extended up to homomorphism $\mu : (V, KG) \rightarrow (V, A)$. Let $\mu(KG) = B \subset A$. Since $(V, A) \in \omega\Sigma$ and (V, B) is a subrepresentation of (V, A) then $(V, B) \in \omega\Sigma$. Hence $(V, KG) \in \omega\Sigma$. Thus $(V, G) \in \eta\Sigma$ and the class $\eta\Sigma$ is a variety of representations of groups. $\square$

Now we describe identities of the variety $\eta\Sigma$. Let $\mathbb{F}$ be a free associative algebra over a field K and let F be a free group.

Definition 4.3 *An element $f \in \mathbb{F}$ is called a polynomial identity of a representation (V, G) if for any homomorphism of algebras $\mu : \mathbb{F} \rightarrow KF$ the element $u = \mu(f) \in KF$ is an identity of the representation (V, G) .*

Remark. It is obvious, that an element $f \in \mathbb{F}$ is a polynomial identity of a representation (V, G) if and only if for any homomorphism $\mu : \mathbb{F} \rightarrow KG$ the element $\mu(f)$ annihilates the module V .

Denote by $I_\Sigma(KF)$ the verbal ideal of KF for the variety Σ . We have the following description of the ideal of identities of the variety $\eta\Sigma$.

Proposition 4.4 ([25]) *The ideal $I_\Sigma(KF)$ is the ideal of identities of the variety $\eta\Sigma$.*

Proof. Let v_i be identities of the variety of associative algebras Σ , $i \in I$. The condition $(V, G) \in \eta\Sigma$ is equivalent to the following one: for any homomorphism $\mu : \mathbb{F} \rightarrow KG$ the elements $\mu(v_i)$ annihilate the module V . Using the remark above we have that the representation (V, G) belongs to $\eta\Sigma$ if and only if all v_i are polynomial identities of (V, G) .

Thus $(V, G) \in \eta\Sigma$ if and only if the images of all v_i in the algebra KF are polynomial identities of the representation (V, G) . These images generate the ideal $I_\Sigma(KF)$. So the class $\eta\Sigma$ is determined by the identities from the ideal $I_\Sigma(KF)$. Since this ideal is invariant under all endomorphisms of the group F then it contains all identities of the class $\eta\Sigma$. $\square$

Remark. The verbal ideal $I_\Sigma(KF)$ of the variety Σ is always invariant under all endomorphisms of the group F , but in this case it is also closed under all endomorphisms of the group algebra KF . However it can be shown that if an ideal in KF is closed under all endomorphisms of the group F this does not mean that this ideal is a homomorphic image of an ideal of identities for some variety of algebras.

Remind that $\mathfrak{M} = \mathfrak{M}(K)$ is the semigroup of varieties of representations of groups and $\mathfrak{M}^a = \mathfrak{M}^a(K)$ is the semigroup of all non-singular varieties of representations of associative algebras over a field K .

Let us consider a map $\eta' : \mathfrak{M}^a \rightarrow \mathfrak{M}$ defined by the rule $\eta'\mathfrak{Y} = \eta\omega^{-1}\mathfrak{Y}$ for every $\mathfrak{Y} \in \mathfrak{M}^a$. We have the following

Proposition 4.5 ([25]) *The map η' is a homomorphism of the semigroup $\mathfrak{M}^a$ to the semigroup $\mathfrak{M}$.*

Proof. Let $\mathfrak{Y}_1$ and $\mathfrak{Y}_2$ be elements of the semigroup $\mathfrak{M}^a$ and let $\mathfrak{Y} = \mathfrak{Y}_1\mathfrak{Y}_2$. Let $\Sigma_1 = \omega^{-1}\mathfrak{Y}_1$, $\Sigma_2 = \omega^{-1}\mathfrak{Y}_2$ and $\Sigma = \omega^{-1}\mathfrak{Y}$. Then we have $\eta'\mathfrak{Y}_1 = \eta\Sigma_1$, $\eta'\mathfrak{Y}_2 = \eta\Sigma_2$ and $\eta'\mathfrak{Y} = \eta\Sigma$.

Let I_1 and I_2 be T -ideals in $\mathbb{F}$ of the varieties Σ_1 and Σ_2 accordingly. Then $I = I_2I_1$ is the T -ideal of the variety Σ . Since the algebra $\mathbb{F}$ and the group F have a countable rank then there is an epimorphism $\mu : \mathbb{F} \rightarrow KF$ and $\mu(I_1) = I_{\Sigma_1}(KF)$, $\mu(I_2) = I_{\Sigma_2}(KF)$, $\mu(I) = I_{\Sigma}(KF)$, moreover

$$\mu(I) = \mu(I_2I_1) = \mu(I_2)\mu(I_1) = I_{\Sigma_2}(KF) \cdot I_{\Sigma_1}(KF).$$

The verbal ideal $I_{\Sigma_2}(KF) \cdot I_{\Sigma_1}(KF)$ determines the variety $\eta\Sigma_1 \cdot \eta\Sigma_2$. Since $I_{\Sigma_2}(KF) \cdot I_{\Sigma_1}(KF) = I_{\Sigma}(KF)$ then the variety $\eta\Sigma_1 \cdot \eta\Sigma_2$ is determined by the ideal $I_{\Sigma}(KF)$. Hence $\eta\Sigma_1 \cdot \eta\Sigma_2 = \eta\Sigma$. $\square$

We can give the following

Definition 4.6 *A variety of representations of groups $\mathfrak{X}$ is called a polynomial variety if there exists a variety of associative algebras Σ such that $\mathfrak{X} = \eta\Sigma$.*

Let us give some examples of polynomial varieties.

Examples.

1. Let Σ be the variety of all commutative algebras over a field K and let $\mathfrak{A}$ be a variety of abelian groups. Then $\omega\mathfrak{A} = \eta\Sigma$.
2. Consider the variety $(\omega\mathfrak{A})^n$. Using the homomorphism $\eta' : \mathfrak{M}^a \rightarrow \mathfrak{M}$ we have $\eta'(\omega\Sigma) = \eta\Sigma = \omega\mathfrak{A}$. Further

$$\eta'((\omega\Sigma)^n) = \eta(\omega^{-1}(\omega\Sigma)^n),$$

$$\eta'((\omega\Sigma)^n) = (\omega\Sigma)^n = (\mathfrak{S} \times \mathfrak{A})^n = \mathfrak{S}^n \times \mathfrak{A},$$

(see section 2.3). A variety of algebras $\Sigma_1 = \omega^{-1}((\omega\Sigma)^n)$ is a variety of algebras with the nilpotent derived ideal of class n . Thus we have $\mathfrak{S}^n \times \mathfrak{A} = \eta\Sigma_1$. So the variety $\mathfrak{S}^n \times \mathfrak{A}$ is polynomial. Remind that the variety $\mathfrak{S}^n$ is a variety generated by the representation $(K^n, UT_n(K))$, and the variety $(\omega\mathfrak{A})^n$ is generated by the representation $(K^n, T_n(K))$.

Proposition 4.5 implies:

Corollary 4.7 ([25]) *All polynomial varieties of representations of groups constitute a subsemigroup in the semigroup $\mathfrak{M}$.* $\square$

Remind that if K is a field then the semigroup $\mathfrak{M}$ of all varieties of representations of groups is free. So we have the following

Problem 4.8 *Is the semigroup of all polynomial varieties free?*

To solve this problem we study some properties of polynomial varieties of representations of groups. In particular, we are looking for interesting examples of polynomial and non-polynomial varieties.

For any representation (V, G) the variety $\widetilde{\text{Var}}(V, G)$ is the variety generated by the representation (V, G) . We denote by $\widetilde{\text{Var}}(V, G)$ the minimal polynomial variety which contains the representation (V, G) , in other words the variety $\widetilde{\text{Var}}(V, G)$ is the minimal polynomial variety containing the variety $\text{Var}(V, G)$. It is not difficult to see that $\widetilde{\text{Var}}(V, G) = \eta\Sigma$, where $\Sigma = \text{Var}(\overline{KG})$ and $\overline{KG} = KG/\text{Ker}(V, KG)$.

For any variety of representations of groups $\mathfrak{X}$ we denote by $\widetilde{\mathfrak{X}}$ the minimal polynomial variety containing the variety $\mathfrak{X}$.

Note that if $\mathfrak{Y} = \text{Var}(V, \overline{KG})$ is a variety of representations of algebras and $\Sigma = \text{Var}(\overline{KG})$ then $\Sigma = \omega^{-1}\mathfrak{Y}$, (remind that $\omega^{-1}\mathfrak{Y} = \overrightarrow{\mathfrak{Y}}$ is the class of all algebras admitting a faithful representation in the variety $\mathfrak{Y}$). Indeed, since the representation $(V, \overline{KG})$ is faithful then $\overline{KG} \in \omega^{-1}\mathfrak{Y}$ and so $\Sigma \subset \omega^{-1}\mathfrak{Y}$. On the other hand, if A is an algebra from $\omega^{-1}\mathfrak{Y}$ then there exists a K -module U such that (U, A) is a faithful representation from the variety $\mathfrak{Y}$. If $y \circ w(x_1, \dots, x_n) \equiv 0$ is an identity in $\mathfrak{Y}$ then $w(x_1, \dots, x_n) \equiv 0$ is an identity of the algebras $\overline{KG}$ and A , since $(V, \overline{KG})$ and (U, A) are faithful representations. So $A \in \Sigma = \text{Var}(\overline{KG})$. Thus $\Sigma = \omega^{-1}\mathfrak{Y}$.

For polynomial varieties we have the property 4.9 which show that the proposition 2.11 is not true for polynomial varieties.

Proposition 4.9 *There are representations (V_1, G_1) and (V_2, G_2) such that*

$$\widetilde{\text{Var}}((V_1, G_1) \nabla (V_2, G_2)) \neq \widetilde{\text{Var}}(V_1, G_1) \cdot \widetilde{\text{Var}}(V_2, G_2).$$

Proof. Let K be a field and let $(V_1, G_1) = (V_2, G_2) = (K, 1)$. Then

$$\widetilde{\text{Var}}((K, 1) \nabla (K, 1)) = \widetilde{\text{Var}}(K^2, \text{UT}_2(K)) = \eta\Sigma,$$

where $\Sigma = \text{Var}A$ and A is the algebra of matrices of the form $\begin{pmatrix} \alpha & \text{End}K \\ 0 & \alpha \end{pmatrix}$, $\alpha \in K$. Note that the variety Σ is the variety of commutative algebras.

In the section 4 we considered the homomorphism $\eta' : \mathfrak{M}^a \rightarrow \mathfrak{M}$ of the semigroup $\mathfrak{M}^a$ of varieties of representations of algebras to the semigroup $\mathfrak{M}$ of varieties of representations of groups. Remind that if $\mathfrak{Y} \in \mathfrak{M}^a$ then $\eta'\mathfrak{Y} = \eta\omega^{-1}\mathfrak{Y}$.

We have the following equalities:

$$\eta'\text{Var}(V, \overline{KG}) = \eta\omega^{-1}\text{Var}(V, \overline{KG}) = \eta\text{Var}(\overline{KG}) = \widetilde{\text{Var}}(V, G).$$

Along with the triangular product of representations of groups we can consider the triangular product of representations of algebras. It is known [8] that for varieties of representations of associative algebras over a field there is the following property:

$$\text{Var}((V_1, A_1) \nabla (V_2, A_2)) = \text{Var}(V_1, A_1) \cdot \text{Var}(V_2, A_2).$$

We denote by $K \cdot 1$ the group algebra over K of the trivial group. So we have

$$\widetilde{\text{Var}}((K, 1)) \cdot \widetilde{\text{Var}}((K, 1)) = \eta'\text{Var}(K, K \cdot 1) \cdot \eta'\text{Var}(K, K \cdot 1) =$$

$$\begin{aligned}
&= \eta' \left(\text{Var}(K, K \cdot 1) \cdot \text{Var}(K, K \cdot 1) \right) = \eta' \left(\text{Var}((K, K \cdot 1) \nabla (K, K \cdot 1)) \right) = \\
&= \eta' \left(\text{Var}(K^2, A_1) \right),
\end{aligned}$$

where A_1 is an algebra of matrices of the form $\begin{pmatrix} K \cdot 1 & \text{End} K \\ 0 & K \cdot 1 \end{pmatrix}$ and this algebra is not commutative.

It is known [25] that a unitriangular product of representations of groups is faithful if and only if all factors are faithful. Since the representation $(K, K \cdot 1)$ is faithful then the representation (K^2, A_1) is also faithful and we have

$$\eta' \left(\text{Var}(K^2, A_1) \right) = \eta \text{Var}(A_1),$$

where $\text{Var}(A_1)$ is not commutative variety. So we have that

$$\widetilde{\text{Var}}((K, 1) \nabla (K, 1)) \neq \widetilde{\text{Var}}(K, 1) \cdot \widetilde{\text{Var}}(K, 1).$$

□

We have following property for the variety $\mathfrak{S}^n$.

Proposition 4.10 *The variety of representations of groups $\mathfrak{S}^n$ is not a polynomial variety.*

Proof. To prove the proposition it is enough to show that the equality $\mathfrak{S}^n = \eta\Sigma$ is impossible for any variety Σ .

Assume that there is a variety of associative algebras Σ such that $\mathfrak{S}^n = \eta\Sigma$. The variety $\mathfrak{S}^n$ contains the representation (K^n, G) , where $G = UT_n(K)$ is the group of unitriangular matrices and the group G acts in a natural way on K^n . The representation (K^n, G) belongs to the variety $\eta\Sigma$ if the linear span $\langle G \rangle_K$ of group G in the algebra $\text{End}(K^n) = M_n(K)$ is an algebra from the variety Σ . Note that the representation (K^n, G) is faithful, so $\overline{G} = G$.

The algebra $\langle G \rangle_K$ is an algebra of matrices of the form:

$$\langle G \rangle_K = \left\{ \begin{pmatrix} \alpha & * & * \\ 0 & \ddots & * \\ 0 & \dots & \alpha \end{pmatrix}, \alpha \in K \right\}$$

Let G_1 be a group of all invertible elements of the algebra $\langle G \rangle_K$. The group G_1 acts faithfully on the module K^n and the linear span $\langle G_1 \rangle_K$ of G_1 in the algebra $\text{End}(K^n)$ coincides with $\langle G \rangle_K$ which is an algebra from the variety Σ . So the representation (K^n, G_1) belongs to the variety $\eta\Sigma = \mathfrak{S}^n$. But it is not true. Thus $\mathfrak{S}^n \neq \eta\Sigma$. □

We have proved that $\mathfrak{S}^n$ is not a polynomial variety. From the proof of the proposition follows that the minimal polynomial variety which contains $\mathfrak{S}^n$ is the variety $\widetilde{\mathfrak{S}^n} = \eta\Sigma$, where Σ is the variety of algebras generated by the algebra of matrices of the following form:

$$\left\{ \begin{pmatrix} \alpha & * & * \\ 0 & \ddots & * \\ 0 & \dots & \alpha \end{pmatrix}, \alpha \in K \right\}$$

Remind that the variety $\omega\mathfrak{A}$ is a polynomial variety (see examples above) and $\omega\mathfrak{A} = \eta\Sigma$, where Σ is a variety of commutative algebras. Moreover, the minimal polynomial variety $\tilde{\mathfrak{S}}$ which contains $\mathfrak{S}$ coincides with $\omega\mathfrak{A}$, that is $\tilde{\mathfrak{S}} = \omega\mathfrak{A}$, since the variety $\mathfrak{S} = \text{Var}(K, 1)$ and the algebra $\langle 1 \rangle_K$ generates the variety of commutative algebras.

Let $\mathfrak{X}$ be a variety of representations of groups over a field K . Let $I_{\mathfrak{X}}(KF) \triangleleft KF$ be the ideal of identities of the variety $\mathfrak{X}$ and $I_{\mathfrak{X}}(KG)$ be the verbal ideal of the group algebra KG . Let Σ be a variety of associative algebras over a field K .

Definition 4.11 Denote by $\Sigma * \mathfrak{X}$ a class of representations of groups over a field K such that a representation (V, G) belongs to $\Sigma * \mathfrak{X}$ if the representation $(V, I_{\mathfrak{X}}(KG))$ lies in $\omega\Sigma$.

In other words, the representation (V, G) belongs to $\Sigma * \mathfrak{X}$ if and only if there is a right ideal J in the algebra KG such that $(V, J) \in \omega\Sigma$ and $(KG/J, G) \in \mathfrak{X}$.

We have

Proposition 4.12 ([25]) For any variety of associative algebras Σ and any variety of representations of groups $\mathfrak{X}$ the class $\Sigma * \mathfrak{X}$ forms a variety of representations of groups. $\square$

Let $\mathbf{N}_n$ be a variety of nilpotent algebras of nilpotency class n (that is $\mathbf{N}_n$ is the variety satisfying the identity $x_1x_2 \dots x_n = 0$). Let $\mathfrak{X}$ be a variety of representations of groups and let $I_{\mathfrak{X}}(KF)$ be the ideal of identities for the variety $\mathfrak{X}$. We have the following

Proposition 4.13 $\mathbf{N}_n * \mathfrak{X} = \mathfrak{X}^n$.

Proof. A representation (V, G) belongs to the variety $\mathbf{N}_n * \mathfrak{X}$ if and only if the algebra $\overline{I_{\mathfrak{X}}(KG)} = I_{\mathfrak{X}}(KG)/\text{Ker}(V, I_{\mathfrak{X}}(KG))$ is an algebra from the variety $\mathbf{N}_n$, that is $(\overline{I_{\mathfrak{X}}(KG)})^n = 0$. So for every $v \in V$ we have $v \circ (\overline{I_{\mathfrak{X}}(KG)})^n = 0$. Since $(I_{\mathfrak{X}}(KG))^n = I_{\mathfrak{X}^n}(KG)$ then $(\overline{I_{\mathfrak{X}}(KG)})^n = \overline{I_{\mathfrak{X}^n}(KG)}$ and $v \circ (\overline{I_{\mathfrak{X}}(KG)})^n = 0$ if and only if $v \circ I_{\mathfrak{X}^n}(KG) = 0$. So the representation $(V, G) \in \mathfrak{X}^n$.

Thus $\mathbf{N}_n * \mathfrak{X} = \mathfrak{X}^n$.

$\square$

Define a PI-variety of representations of groups

Definition 4.14 A variety of representations of groups $\mathfrak{X}$ is called a PI-variety if every representation from $\mathfrak{X}$ satisfies some non-trivial polynomial identity.

There is the following

Proposition 4.15 ([25]) If a variety $\mathfrak{X}$ is generated by a representation (V, G) satisfying a polynomial identity f then all representations from $\mathfrak{X}$ satisfy this identity.

Proof. Let $\mathfrak{X}$ be a variety generated by the representation (V, G) and let (V, G) satisfies a polynomial identity f . Let Σ be a variety of algebras determined by the identity f . Then (V, G) belongs to a variety $\eta\Sigma$. Hence $\text{Var}(V, G) = \mathfrak{X}$ is contained in the variety $\eta\Sigma$. So all representations from $\mathfrak{X}$ satisfy the polynomial identity f . $\square$

Corollary 4.16 ([25]) *If $\mathfrak{X}$ is a polynomial variety then there is a non-trivial polynomial identity such that all representations from $\mathfrak{X}$ satisfy this identity.*
 $\square$

Remark. Let $\mathbb{F}$ be a free associative algebra over a field K with free generates $z_1, z_2, \dots$ and let F be a free group with free generates $x_1, x_2, \dots$. Now let $f = f(z_1, \dots, z_n) \in \mathbb{F}$ be a multilinear polynomial. Consider a map $\mathbb{F} \rightarrow F$ by the rule $z_i \rightarrow x_i$. The polynomial $f = f(z_1, \dots, z_n)$ is an identity of a representation (V, G) if and only if the representation (V, G) satisfies the identity $f' = f(x_1, \dots, x_n)$.

It is well known that every PI -algebra satisfies a multilinear identity. Hence every PI -variety of representations of groups satisfies some non-trivial multilinear identity. The remark above allows us to consider this identity as an element of the group algebra KF .

If K is a field of zero characteristic then every polynomial identity are equivalent to some multilinear identity. Thus every polynomial varieties of representations of groups over a field of zero characteristic can be determined by the multilinear identities.

5 General view on dimension subgroups

Recall that the classical dimension subgroup $D_n(\mathbb{Z}, G)$ is a subgroup of the group G over $\mathbb{Z}$ is defined as follows: $D_n(\mathbb{Z}, G)$ is the set of all elements $g \in G$ such that $(g - 1)$ belongs to the ideal $(\Delta_G)^n \subset \mathbb{Z}G$, where Δ_G is the augmentation ideal of $\mathbb{Z}G$. That is

$$D_n(K, G) = (1 + (\Delta_G)^n) \cap G.$$

The exposition of the classical dimension subgroup, various problems concerning dimension subgroups and book review are contained in the book of R. Mikhailov and I.B.S. Passi [16]. In particular, we would note the papers by W. Magnus [12], O. Grün [6], N. Gupta [7], I.B.S. Passi [19], E. Rips [26], E. Witt [30]. We are interested to consider the dimension subgroup from more general point of view. Remind that the n -th power $(\Delta_G)^n \subset KG$ of the augmentation ideal (Δ_G) is the ideal of identities of the variety of representations of groups $\mathfrak{S}^n$. So this insight yields a new approach to dimension subgroups.

5.1 Dimension subgroup for a variety of representations of groups

Let $\mathfrak{X}$ be a variety of representations of groups. We consider representations (V, G) of the group G contained in the variety $\mathfrak{X}$. We define a subgroup $D_{\mathfrak{X}}(G)$ of the group G as follows:

Definition 5.1 *A dimension subgroup $D_{\mathfrak{X}}(G)$ of a group G for a variety of representations of groups $\mathfrak{X}$ is the intersection of kernels of all representations $(V, G) \in \mathfrak{X}$.*

Denote by $\overrightarrow{\mathfrak{X}}$ the class of groups admitting faithful representation in the variety $\mathfrak{X}$. This class is a quasi-variety of groups and the corresponding verbal

subgroup coincides with the dimension subgroup. In particular, if K is a field of zero characteristic then $\mathfrak{S}^{\vec{n}}$ is the quasi-variety of all torsion free nilpotent groups of the class $(n-1)$ [14]. For the quasi-variety $\mathfrak{S}^{\vec{n}}$ there is also the problem of B. Plotkin [25] which asks: has the quasi-variety $\mathfrak{S}^{\vec{n}}$ a finite basis of quasi-identities? For the quasi-variety $\mathfrak{S}^{\vec{1}}$ this problem is solved by R. Mikhailov and I.B.S. Passi [17] (see also [16]). They proved that this quasi-variety is not finitely based.

Let $I_{\mathfrak{X}}(KF)$ be the ideal of identities of the group algebra KF for a variety $\mathfrak{X}$ (here F is a free group) and let $I_{\mathfrak{X}}(KG)$ be the verbal ideal of KG for the variety $\mathfrak{X}$. The representation $(KG/I_{\mathfrak{X}}(KG), G)$ is the free representation of the group G in the variety $\mathfrak{X}$. The kernel of this representation is a normal subgroup $H = (1 + I_{\mathfrak{X}}(KG)) \cap G$ consisting of all $g \in G$ such that the element $(g-1)$ belongs to the ideal $I_{\mathfrak{X}}(KG)$.

We have the following proposition.

Proposition 5.2 ([25]) *The kernel of the free representation $(KG/I_{\mathfrak{X}}(KG), G)$ coincides with the dimension subgroup $D_{\mathfrak{X}}(G)$ of the group G for the variety $\mathfrak{X}$.*

Proof. By the definition of $D_{\mathfrak{X}}(G)$ we have $D_{\mathfrak{X}}(G) \subseteq H$, where $H = (1 + I_{\mathfrak{X}}(KG)) \cap G$ is the kernel of the free representation $(KG/I_{\mathfrak{X}}(KG), G)$. We will show that $H \subseteq D_{\mathfrak{X}}(G)$.

Let (V, G) be a representation of the group G such that $(V, G) \in \mathfrak{X}$, let v be an element in V . The map $g \rightarrow v \circ g$ induces the homomorphism of modules: $KG \rightarrow V$, and this homomorphism is permutable with the action of the group G . Since the ideal $I_{\mathfrak{X}}(KG)$ annihilates the module V then there is a homomorphism $\nu : KG/I_{\mathfrak{X}}(KG) \rightarrow V$. Let g be an arbitrary element from the kernel H and let $\bar{1} = 1 + I_{\mathfrak{X}}(G)$. Then for all $v \in V$ we have:

$$v \circ g = (\bar{1})^\nu \circ g = (\bar{1} \circ g)^\nu = (\bar{1})^\nu = v.$$

So any element $g \in H$ belongs to the kernel of the representation (V, G) . It means that the kernel of the representation $(KG/I_{\mathfrak{X}}(KG), G)$ is contained in the kernel of every representation (V, G) form the variety $\mathfrak{X}$. Thus $D_{\mathfrak{X}}(G)$ coincides with the kernel of the free representation $(KG/I_{\mathfrak{X}}(KG), G)$. $\square$

Since the ideal $(\Delta_G)^n$ is the ideal of identities of the variety $\mathfrak{S}^n$ the classical dimension subgroup $D_n(K, G)$ is the dimension subgroup of the group G for the variety $\mathfrak{X} = \mathfrak{S}^n$, that is $D_n(K, G) = D_{\mathfrak{S}^n}(G)$. So we have that the classical dimension subgroup connects with the dimension subgroup for the variety of representations of groups $\mathfrak{S}^n$.

It is interesting to consider the following

Problem 5.3 *Describe the dimension subgroups for the polynomial varieties $\mathfrak{S}^n$ and $(\mathfrak{S})^n = (\omega\mathfrak{A})^n$, respectively.*

We have a partial solution of this problem for the free group F and the variety $\widetilde{\mathfrak{S}}^n$, see Theorem 5.10 below.

Remind that the variety $\mathfrak{S}^n$ is the variety generated by the representation $(K^n, UT_n(K))$, and the variety $(\mathfrak{S})^n = (\omega\mathfrak{A})^n$ is generated by the representation $(K^n, T_n(K))$.

In the section 5.3 we consider the concept of a dimension subgroup for variety of associative algebras and some relations between these subgroups.

5.2 Linearization of groups and *PI*-groups

5.2.1 Let A be an associative algebra with unit over a field K and let G be a group.

Definition 5.4 An arbitrary homomorphism $\rho: G \rightarrow A$ of a group G into the group of invertible elements of an algebra A is called a linearization of the group G in the algebra A over a field K .

For a given group G we consider the category of all K -linearizations of G over a field K . The objects of this category are linearizations $\rho_i: G \rightarrow A_i$ of the group G in the K -algebras A_i . Let ρ_1 and ρ_2 be two linearizations of the group G in the algebras A_1 and A_2 , accordingly. Then there is a homomorphism μ of the algebra A_1 to the algebra A_2 such that

$$\begin{array}{ccc} & \nearrow \rho_1 & A_1 \\ G & & \downarrow \mu \\ & \searrow \rho_2 & A_2 \end{array}$$

So the morphisms of this category are commutative diagrams above.

This category possesses a universal initial object. It is identical embedding $G \xrightarrow{id} KG$ of the group G into the group algebra KG . For every linearization $\rho: G \rightarrow A$ we have a homomorphism $\mu: KG \rightarrow A$ such that the following diagram is commutative

$$\begin{array}{ccc} G & \xrightarrow{id} & KG \\ & \searrow \rho & \downarrow \mu \\ & & A \end{array}$$

Definition 5.5 A linearization $\rho: G \rightarrow A$ is called faithful if the map ρ is an injective homomorphism.

The linearization $G \rightarrow KG$ is faithful. In general case, if $\rho: G \rightarrow A$ is a linearization of a group G and $G^\rho = G/\text{Ker}\rho$ then the linearization $\bar{\rho}: G^\rho \rightarrow A$ is faithful.

If a linearization $\rho: G \rightarrow A$ is a faithful linearization then the group G can be considered as a subgroup of the group of invertible elements of the algebra A .

Let Σ be a non-trivial variety of associative algebras over a fixed field K . We will consider a linearization $\rho: G \rightarrow A$, where A is a K -algebra from the variety Σ .

We also have the category of such linearizations and its initial object is a linearization $\rho: G \rightarrow KG/I_\Sigma(KG)$, where $I_\Sigma(KG)$ is the verbal ideal in KG for the variety Σ . This linearization can be not faithful.

Definition 5.6 A group G is called a *PI*-group if there is a faithful linearization $\rho: G \rightarrow A$, where A is an algebra from some non-trivial variety of associative algebras Σ .

In more detail we consider *PI*-group in [1] where we solve for these groups some Burnside-type problems.

Note that, if a group G admits a faithful linearization in an algebra A from a variety Σ then the linearization $G \rightarrow KG/I_\Sigma(KG)$ is also faithful.

5.2.2 Let us return to dimension subgroups.

Let $\rho : G \rightarrow A$ be a linearization of a group G in an algebra A from a variety Σ . In particular we can consider universal linearization $G \rightarrow KG/I_\Sigma(KG)$, where $I_\Sigma(KG)$ is the verbal ideal in KG of the variety Σ . It is clear that the kernel of every homomorphism $\rho : G \rightarrow A$ contains the kernel of universal linearization. Observe that the kernel of the universal linearization is the set of all $g \in G$ such that the element $(g - 1) \in KG$ belongs to the ideal $I_\Sigma(KG)$. This kernel is denoted by $D_\Sigma(G)$. So we have

$$D_\Sigma(G) = \{ g \in G \mid g - 1 \in I_\Sigma(KG) \}.$$

Definition 5.7 *The kernel $D_\Sigma(G)$ of universal linearization $G \rightarrow KG/I_\Sigma(KG)$ is called a dimension subgroup of the group G for the variety of algebras Σ .*

Note that a group G is a *PI*-group with given variety Σ if and only if $D_\Sigma(G) = 1$.

Let $\mathfrak{D}(\Sigma)$ be a class of all *PI*-groups with the given Σ . Here we have a theorem similar to the theorem for the varieties of representations of groups. The latter one states that a class of all groups G admitting a faithful representation (V, G) from some variety of representations of groups $\mathfrak{X}$ forms a quasivariety of groups [25].

Theorem 5.8 *The class $\mathfrak{D}(\Sigma)$ is a quasivariety of groups.*

Proof. To prove the theorem it is sufficient to show [13] that the class $\mathfrak{D}(\Sigma)$ contains the trivial group and is closed under subgroups and filtered products. It is obvious that for class $\mathfrak{D}(\Sigma)$ first two conditions hold true. We need to prove that the class $\mathfrak{D}(\Sigma)$ is closed under the filtered products.

Let I be a non-empty set, let D be a filter on the set I . Let G_α be groups from the class $\mathfrak{D}(\Sigma)$, let $\varphi_\alpha : G_\alpha \rightarrow A_\alpha$ be faithful linearizations of groups G_α in the algebras A_α from the variety Σ .

Let $A = \prod_\alpha A_\alpha$ be the Cartesian product of the algebras A_α . Let A/J be the filtered product of these algebras, where J is the ideal of A such that

$$a \in J \Leftrightarrow \{ \alpha \in I \mid a(\alpha) = 0 \} \subset D.$$

Since all A_α are *PI*-algebras from the variety Σ then the algebra A and the algebra A/J belongs to the variety Σ .

Let $G = \prod_\alpha G_\alpha$ be the Cartesian product of the groups G_α . Let G/H be the filtered product of the groups G_α , where H is the subgroup of G such that

$$h \in H \Leftrightarrow \{ \alpha \in I \mid h(\alpha) = 1 \} \subset D.$$

Consider the linearization $\psi : G \rightarrow A/J$ defined by the rule $f \rightarrow f' + J$ where $f'(\alpha) = \varphi_\alpha(f(\alpha))$ for all $\alpha \in I$, $f \in G$. It is easy to see that ψ is a homomorphism.

Consider the kernel of the homomorphism ψ :

$$\text{Ker}(\psi) = \{ g \mid \psi(g) = 1 + J \}.$$

For any element $g \in \text{Ker}(\psi)$ we have $\psi(g)(\alpha) = \varphi_\alpha(g(\alpha)) + J(\alpha) = 1(\alpha) + J(\alpha) = 1(\alpha)$, so $\varphi_\alpha(g(\alpha)) = 1(\alpha)$ and $g \in H$. Thus $\text{Ker}(\psi) = H$.

Then $G/H \cong \text{Im } \psi \subset A/I$ and the group G/H admits the faithful linearization in the algebras from the variety Σ . Thus the class $\mathfrak{D}(\Sigma)$ is closed under the filtered products.

So the class $\mathfrak{D}(\Sigma)$ is a quasivariety. $\square$

5.3 Dimension subgroup for varieties of representations of groups and for varieties of associative algebras

5.3.1 Now we will compare two approaches (see Definition 5.1 and Definition 5.7) to the notion of a dimension subgroup.

Let $\mathfrak{X}$ be a variety of representations of groups and let $I_{\mathfrak{X}}(KG)$ be the verbal ideal in KG for the variety $\mathfrak{X}$. Let Σ be a variety of associative algebras over a field K and let $I_{\Sigma}(KG)$ be the verbal ideal in KG for the variety Σ . In the section 4 (see Proposition 4.4) it was proved that if $\mathfrak{X} = \eta\Sigma$ then $I_{\mathfrak{X}}(KF) = I_{\Sigma}(KF)$, where F is the free group.

Hence, here we have the following

Theorem 5.9 *If $\mathfrak{X} = \eta\Sigma$ is a polynomial variety then the dimension subgroup $D_{\mathfrak{X}}(G)$ coincides with the dimension subgroup $D_{\Sigma}(G)$.* $\square$

Remark. Remind that there is an equality $\mathfrak{S}^n \times \mathfrak{A} = (\omega\mathfrak{A})^n = \eta\Sigma$ (see section 4), where $\mathfrak{A}$ is the variety of abelian groups and Σ is the variety of associative algebras satisfying the identity

$$(x_1y_1 - y_1x_1) \dots (x_ny_n - y_nx_n) \equiv 0.$$

So, we have the following equality

$$D_{\mathfrak{S}^n \times \mathfrak{A}} = D_{\Sigma}.$$

Remind that $D_n(K, G) = (1 + (\Delta_G)^n) \cap G$ is the dimension subgroup of a group G over a ring K , where Δ_G is the augmentation ideal of KG , and for every group G the dimension subgroup $D_n(K, G)$ coincides with the dimension subgroup $D_{\mathfrak{S}^n}(G)$ for the variety of representations of groups $\mathfrak{S}^n$. For a free group F we have the following

Theorem 5.10 *For a free group F its dimension subgroup $D_n(K, F)$ coincides with the dimension subgroup $D_{\widetilde{\mathfrak{S}^n}}(F)$ of the group F for the polynomial variety of representations of groups $\widetilde{\mathfrak{S}^n}$.*

Proof.

In the section 4 it was shown that $\widetilde{\mathfrak{S}^n} = \eta\Sigma$, where Σ is a variety of algebras generated by the algebra of matrices of the following from:

$$\left\{ \begin{pmatrix} \alpha & * & * \\ 0 & \ddots & * \\ 0 & \dots & \alpha \end{pmatrix}, \alpha \in K \right\},$$

and this algebra is the linear span $\langle G \rangle_K$ of the group $G = UT_n(K)$ of unitriangular matrices in the algebra $\text{End}(K^n)$.

Let $H = (\langle G \rangle_K)^*$ be the group of invertible elements of the algebra $\langle G \rangle_K$, that is

$$H = \left\{ \begin{pmatrix} \alpha & * & * \\ 0 & \ddots & * \\ 0 & \dots & \alpha \end{pmatrix} \mid \alpha \in K, \alpha \neq 0 \right\}.$$

The group $H = UT_n(K) \cdot Z(T_n(K)) = UT_n(K) \times Z(T_n(K))$, where $Z(T_n(K))$ is the center of the group $T_n(K)$. The group $UT_n(K)$ is nilpotent of the nilpotency class $(n-1)$ and the group $Z(T_n(K))$ is abelian hence the group H is nilpotent of the class $(n-1)$.

Let F be a free group. Consider all possible homomorphisms of the group F to the group H . These homomorphisms induce homomorphisms of the group algebra KF to the group algebra $KH = \langle G \rangle_K$:

$$\psi_i : KF \rightarrow \langle G \rangle_K.$$

Let U_i be the kernel of the homomorphism ψ_i and let $U = \bigcap_i U_i$ be the intersection of kernels of all possible homomorphisms ψ_i . So U is the set of all identities of the algebra $\langle G \rangle_K$ in the algebra KF , in other words, U is the verbal ideal of the variety Σ in the group algebra KF :

$$U = I_\Sigma(KF).$$

Using the proposition 4.4 we have:

$$I_\Sigma(KF) = I_{\eta\Sigma}(KF) = I_{\widetilde{\mathfrak{S}^n}}(KF).$$

According to the Remak's theorem the algebra KF/U is a subalgebra of the Cartesian product of KF/U_i . All algebras KF/U_i are subalgebras of $\langle G \rangle_K$. So the algebra KF/U can be embedded into a Cartesian power of the algebra $\langle G \rangle_K$. Denote this cartesian power by $\prod \langle G \rangle_K$.

Then, $\prod H$ is the group of all invertible elements of the algebra $\prod \langle G \rangle_K$. So the group $(KF/U)^*$ of all invertible elements of the algebra KF/U is a subgroup of $\prod H$.

The group H is nilpotent of the class $(n-1)$. Thus, the group $\prod H$ is also nilpotent of the class $(n-1)$. Hence the group $(KF/U)^*$ is nilpotent of nilpotency class at most $(n-1)$.

Let $D_{\widetilde{\mathfrak{S}^n}}(F)$ be the dimension subgroup of the free group F for the variety $\widetilde{\mathfrak{S}^n}$, that is

$$D_{\widetilde{\mathfrak{S}^n}}(F) = \{g \in F \mid (g-1) \in U = I_{\widetilde{\mathfrak{S}^n}}(KF)\}.$$

Since $I_{\widetilde{\mathfrak{S}^n}}(KF) = I_\Sigma(KF)$ then $D_{\widetilde{\mathfrak{S}^n}}(F) = D_\Sigma(F)$ and the subgroup $D_\Sigma(F)$ is a kernel of the homomorphism $F \rightarrow KF/U$. Then we have

$$F/D_{\widetilde{\mathfrak{S}^n}}(F) \subset (KF/U)^*.$$

It means that the group $F/D_{\widetilde{\mathfrak{S}^n}}(F)$ is nilpotent of the class nilpotency at most $(n-1)$. Let $\gamma_n(F)$ is the n -th term of the lower central series of the group F . So we have

$$\gamma_n(F/D_{\widetilde{\mathfrak{S}^n}}(F)) = 1,$$

and then

$$\gamma_n(F) \subset D_{\widetilde{\mathfrak{S}^n}}(F).$$

It is known a fundamental result that for a free group F there is the equality $\gamma_n(F) = D_n(\mathbb{Z}, F)$ [12], [6], [30] (see also [27], [16]). So for a field K of zero characteristic we have

$$\gamma_n(F) = D_n(K, F).$$

Thus we have

$$D_n(K, F) \subset D_{\widetilde{\mathfrak{S}^n}}(F).$$

On the other hand, $\mathfrak{S}^n \subset \widetilde{\mathfrak{S}^n}$ and then $I_{\widetilde{\mathfrak{S}^n}}(KF) \subset I_{\mathfrak{S}^n}(KF)$, where $I_{\widetilde{\mathfrak{S}^n}}(KF)$ and $I_{\mathfrak{S}^n}(KF)$ are verbal ideals of KF for the varieties $\widetilde{\mathfrak{S}^n}$ and $\mathfrak{S}^n$, respectively. Hence

$$D_{\widetilde{\mathfrak{S}^n}}(F) \subset D_{\mathfrak{S}^n}(F) = D_n(K, F).$$

Thus

$$D_n(K, F) = D_{\widetilde{\mathfrak{S}^n}}(F).$$

Theorem is proved. $\square$

We have the following

Corollary 5.11 *For a free group F the dimension subgroup $D_{\mathfrak{S}^n}(F)$ for the variety of representations of groups $\mathfrak{S}^n$ coincides with the dimension subgroup $D_{\widetilde{\mathfrak{S}^n}}(F)$ for the polynomial variety $\widetilde{\mathfrak{S}^n}$.*

We can state

Problem 5.12 *Find a non-free group G such that the dimension subgroup $D_n(K, G)$ coincides with the dimension subgroup $D_{\mathfrak{X}}(G)$ for some variety of representations of groups $\mathfrak{X} \neq \mathfrak{S}^n$.*

5.3.2 Let $\mathfrak{X} = \eta\Sigma$ be a polynomial variety of representations of groups. Assume that the identities of the variety Σ are known.

Let G be a group, let KG be the group algebra of G and let $I_{\mathfrak{X}}(KG)$ be the verbal ideal of identities for the variety $\mathfrak{X}$ in the algebra KG . Note that we have the equality $I_{\mathfrak{X}}(KG) = I_{\Sigma}(KG)$, where $I_{\Sigma}(KG)$ is the verbal ideal of identities for the variety of algebras Σ in the algebra KG . The group G admits the representation $\rho = (KG/I_{\mathfrak{X}}(KG), G)$ from the variety $\mathfrak{X}$ and the kernel of this representation is the dimension subgroup of G for the variety $\mathfrak{X}$, that is

$$D_{\mathfrak{X}}(G) = \text{Ker}\rho = \{g \in G \mid g - 1 \in I_{\mathfrak{X}}(KG)\}.$$

Moreover we have (see theorem 5.9):

$$D_{\mathfrak{X}}(G) = D_{\Sigma}(G).$$

Along with the representation $\rho = (KG/I_{\mathfrak{X}}(KG), G)$ we can consider the representation $\rho_1 = (KG/I_{\mathfrak{X}}(KG), KG)$. Since the representation ρ belongs to the variety $\mathfrak{X} = \eta\Sigma$ then the algebra $\overline{KG} = KG/\text{Ker}\rho_1 \cong \langle \overline{G} \rangle$ is an algebra from the variety Σ , (where $\langle \overline{G} \rangle$ is a linear span of the group $\overline{G} = G/\text{Ker}\rho = G/D_{\Sigma}(G)$ in the algebra $\text{End}(KG/I_{\mathfrak{X}}(KG))$). Thus, $\overline{G} \subset \overline{KG}$.

We have the following

Problem 5.13 We know the identities of the algebra $\overline{KG} \in \Sigma$. What can we say about the identities of the group $\overline{G} = G/D_\Sigma(G)$?

We have solution of this problem for the variety Σ of algebras with the nilpotent derived ideal.

Let $\mathfrak{N}_{n-1}$ be the variety of nilpotent groups of class $(n-1)$, and let $\mathfrak{A}$ be a variety of abelian groups.

Proposition 5.14 *If Σ is a variety of algebras with the nilpotent derived ideal of class n and a group G admits a representation from the variety $\eta\Sigma$ then the group $G/D_\Sigma(G)$ belongs to the variety of groups $\mathfrak{N}_{n-1} \cdot \mathfrak{A}$.*

Proof. Let $\mathfrak{X} = \eta\Sigma$. In the section 4 it was showed that $\eta\Sigma = \mathfrak{S}^n \times \mathfrak{A} = (\omega\mathfrak{A})^n$.

Let G be a group and let $V = KG/I_\Sigma(KG)$ then there is the representation (V, G) from the variety $\mathfrak{X}$. Let $\overline{G} = G/D_\Sigma(G)$. Since the representation $(V, \overline{G})$ belongs to the variety $\mathfrak{X} = \mathfrak{S}^n \times \mathfrak{A}$ then there is a normal subgroup $\overline{H} \triangleleft \overline{G}$ such that there is a representation $(V, \overline{H})$ from the variety $\mathfrak{S}^n$ and the group $\overline{G}/\overline{H}$ is abelian.

The representation $(V, \overline{H})$ is faithful (as a subrepresentation of the faithful representation $(V, \overline{G})$) and it belongs to the variety $\mathfrak{S}^n$, so according to the Kaloujnine's theorem [9] the group $\overline{H}$ is $(n-1)$ -nilpotent. Moreover $\overline{G}/\overline{H}$ is abelian and the group $\overline{G}$ belongs to the variety $\mathfrak{N}_{n-1} \cdot \mathfrak{A}$. $\square$

Let $\overline{G} = G/D_\Sigma(G)$ be a group from the variety of groups Θ . Let B be the verbal subgroup of the group G for the variety Θ . Then $G/B \in \Theta$ and $B \subset D_\Sigma(G)$.

We can consider

Problem 5.15 *What can we say about the structure of the group $D_\Sigma(G)/B$?*

Let B be the verbal subgroup of the group G for the variety $\mathfrak{N}_{n-1} \cdot \mathfrak{A}$. Let Σ be the variety of algebras with the nilpotent derived ideal of the class n . We have the following

Proposition 5.16 *Let a group G admit a representation in the variety of representations of groups $\eta\Sigma$. If the group G/B is torsion-free then $D_\Sigma(G) = B$.*

Proof. As it was noted above $B \subset D_\Sigma(G)$. We will show that $D_\Sigma(G) \subset B$.

Let $G_1 = G/B$. Since $B \subset D_\Sigma(G)$ and $G/D_\Sigma(G) \in \mathfrak{N}_{n-1} \cdot \mathfrak{A}$ then the group $G_1 \in \mathfrak{N}_{n-1} \cdot \mathfrak{A}$. So there is a normal subgroup $G_2 \triangleleft G_1$ such that G_2 is $(n-1)$ -nilpotent and the group G_1/G_2 is abelian. Since the group G_1 is torsion-free then the subgroup G_2 is also torsion-free and $(n-1)$ -nilpotent. According to the result of Maltsev [14] (see also [25]) if K is a field of a characteristic zero then the class of all groups admitting a faithful representation in the variety $\mathfrak{S}^n$ coincides with the class $\mathfrak{N}_{n-1,0}$ of all torsion-free nilpotent groups. So there is a K -module U such that the group G_2 admits a faithful representation (U, G_2) from the variety $\mathfrak{S}^n$.

So we have the group G_1 and the representation (U, G_2) of the subgroup G_2 of the group G_1 . Here we can use induced representations. Let $U_1 = U \otimes_{KG_2} KG_1$,

let (U_1, G_1) be a representation induced by the representation (U, G_2) . So for every $u \in U_1$ and every $g \in G_1$ we have

$$u_1 \circ g = (u \otimes \sum_i \alpha_i g_i) \circ g = (u \circ g) \otimes \sum_i \alpha_i g_i = u \otimes \sum_i \alpha_i g g_i,$$

where $u \in U$, $\alpha_i \in K$, $g_i \in G_1$. Since the representation (U, G_2) is faithful the representation (U_1, G_1) is also faithful. Let $h \in G_2$ then

$$(U \otimes_{KG_2} KG_1) \circ h = (U \circ h) \otimes_{KG_2} KG_1.$$

The group G_2 acts n -unitriangularly on U . Then G_2 acts n -unitriangularly on $U \otimes_{KG_2} KG_1$ and the representation (U_1, G_2) belongs to the variety $\mathfrak{S}^n$.

Thus we have that the group G_1 has the normal subgroup G_2 such that G_1/G_2 is abelian and $(U_1, G_2) \in \mathfrak{S}^n$, it means that the faithful representation (U_1, G_1) belongs to the variety $\mathfrak{S}^n \times \mathfrak{A}$.

With the representation $(U_1, G_1) = (U_1, G/B)$ we consider the representation (U_1, G) . It belongs to the variety $\mathfrak{S}^n \times \mathfrak{A}$ and the kernel of this representation is the subgroup B . Since $D_\Sigma(G) = D_{\mathfrak{X}}(G)$ is the intersection of kernels of all representations of the group G from the variety $\mathfrak{X} = \eta\Sigma$ then $D_\Sigma(G) \subset B$.

So we have proved that $D_\Sigma(G) = B$. $\square$

Remind that the variety $(\omega\mathfrak{A})^n$ is generated by the representation $(K^n, T_n(K))$ and the variety $\mathfrak{S}^n$ is generated by the representation $(K^n, UT_n(K))$. So the classical dimension subgroup is related to the group of unitriangular matrices $UT_n(K)$ and dimension subgroup for the polynomial variety $(\omega\mathfrak{A})^n$ is related to the group of triangular matrices $T_n(K)$.

References

- [1] E.V. Aladova, B.I. Plotkin, *PI*-groups and *PI*-representations of groups. Fundamental and applied mathematics, 2009, to appear. Arxiv: math RA: 0905.3651.
- [2] G.M. Bergman, J. Lewin, The semigroup of ideals of a fir is (usually) free, J. London Math. Soc., **11**, (1975), no. 1, p. 21–31.
- [3] G. Birkhoff, On the structure of abstract algebras, Proc. Cambridge Philos. Soc., **31**, (1935), p. 433–454.
- [4] R. Fox, Free differential calculus. I. Derivation in the free group ring. Ann. of Math., **57**, (1953), no. 2, p. 547–560.
- [5] A.C. Grinberg, Varieties of stable linear representations. (Russian). Latvian math. yearbook. Izdat. "Zinatne", Riga, **9**, (1971), p. 39–46.
- [6] O. Grün, Über die Faktogruppen freier Gruppen I. Deutsche Math. (Jahrgang 1), **6**, (1936), p. 772–782.
- [7] N. Gupta, The dimension subgroup conjecture. Bull. London Math. Soc., **22**, (1990), no. 5, p. 453–456.
- [8] U.E. Kaljulaid, Triangular products of representations of semigroups and associative algebras, Uspehi Math. Nauk, **32**, no. 4, (1977), p. 253–254.

- [9] L. Kaloujnine, Über gewisse Beziehungen zwischen einer Gruppe und ihren Automorphismen. (German), Berliner Math. Tagung, (1953), pp. 164–172.
- [10] L.E. Krop, Solvable varieties of pairs. (Russian) Latvian. Mat. yearbook. **18**, (1976), p. 64–80.
- [11] S. Mac Lane, Categories for the working mathematician, Second edition. Graduate text in mathematics, 5. Springer-Verlag, New York, (1998), xii+314 pp.
- [12] W. Magnus, Über Beziehungen zwischen höheren Kommutatoren. J. reine angew. Math., **177**, (1937), p. 105–115.
- [13] A.I. Malcev, Algebraic systems. Posthumous edition, edited by D. Smirnov and M. Taiclin. Translated from the Russian by B. D. Seckler and A. P. Doohovskoy. Die Grundlehren der mathematischen Wissenschaften, Band 192. Springer-Verlag, New York-Heidelberg, 1973.
- [14] A.I. Malcev, Generalized nilpotent algebras and their associated groups. (Russian) Mat. Sbornik N.S. 25(67), (1949). 347–366.
- [15] R. Mikhailov, I.B.S. Passi, Augmentation powers and group homology. J. Pure Appl. Algebra **192** (2004), no. 1-3, p. 225–238.
- [16] R. Mikhailov, I.B.S. Passi, Lower central and dimension series of groups. Lecture Notes in Mathematics, 1952. Springer-Verlag, Berlin, 2009. xxii+346 pp.
- [17] R. Mikhailov, I.B.S. Passi, The quasi-variety of groups with trivial fourth dimension subgroup. J. Group Theory, **9**, (2006), p. 369–381.
- [18] B.H. Neumann, H. Neumann, P.M. Neumann, Wreath products and varieties of groups, Math. Ztschr., **80**, (1962), no. 1, p. 44–62.
- [19] I.B.S. Passi, Group rings and their augmentation ideals. Lecture Notes in Mathematics, **715**. Springer, Berlin, (1979). vi+137 pp.
- [20] B.I. Plotkin, Group of automorphisms of algebraic system. Translated from the Russian by K.A. Hirsh. Wolters-Noordhoff Publishing, Groningen, (1972), xviii+502 pp.
- [21] B.I. Plotkin, Notes on Engel groups and Engel elements in groups. Some generalizations, (English, Russian summary), Izv. Ural. Gos. Univ. Mat. Mekh., **36**, (2005), no. 7, p. 153–166, p. 192–193.
- [22] B.I. Plotkin, Varieties of groups, and varieties of pairs that are connected with representations of groups. (Russian), Sibirsk. Mat. Z. **13**, (1972), no. 5, p. 1030–1053.
- [23] B.I. Plotkin, The varieties and quasivarieties that are connected with group representations, (Russian), Dokl. Akad. Nauk SSSR, **196** (1971), no. 3, p. 527–530.
- [24] B.I. Plotkin, A.S. Grinberg, Semigroups of varieties that are connected with group representations. (Russian), Sibirsk. Mat. Z., **13**, (1972), no. 5, p. 841–858.

- [25] B.I. Plotkin, S.M. Vovsi, Varieties of group representations. General theory, connections and applications, (Russian), "Zinatne", Riga, (1983), 339 pp.
- [26] E. Rips, On the fourth integer dimension subgroup. Israel J. Math., **12**, (1972), p. 342–346.
- [27] F. Röhl, Review and some critical comments on a paper of Grun concerning the dimension subgroup conjecture. Bol. Soc. Brasil. Mat. **16**, no. 2, (1985), p. 11–27.
- [28] A.L. Smel'kin, Semigroup of group varieties, (Russian), Dokl. Akad. Nauk SSSR, **149**, (1963), no. 3, p. 543–545.
- [29] S.M. Vovsi, Triangular products of group representations and their applications. Progress in Mathematics, 17. Birkhauser, Boston, Mass., 1981. ix+127 pp.
- [30] E. Witt, Treue Darstellung Lieschen Ringe. J. Reine angew. Math., **177**, (1937), p. 152–160.